

## Sulphur cinquefoil

### *Potentilla recta*

#### Description

Used for its medicinal properties. Also fruit of plant is edible.

#### Habit

Erect; perennial; 16-32 in tall; hairy forb; unbranched to the inflorescence; woody taproot with short branch roots, no rhizomes.

#### Leaves


Palmately-divided, deeply-toothed, lower leaves long-stalked with 5-7 leaflets; upper leaves smaller, shorter stalked and with 3 leaflets.

#### Stems

Erect; upright; slender; 20-80 cm (8-30 in) tall; upper portion may be branched; long hairs perpendicular to leafstalk and stem.

#### Flowers

Pale yellow in color, 5-parted, 3/4 in wide, inflorescence of many flowers in flattened, branched cluster (cyme); blooms June through





August.

### **Fruits and Seeds**

Achene strongly veined, brownish-purple, 1 mm long.

### **Habitat**

Native to Eurasia. Found in dry, disturbed sites.

### **Reproduction**

By seed and vegetatively by sprouting from a caudex.

### **Similar**

Foliage has some resemblance to Marijuana (*Cannabis sativa*) but marijuana is taller with different flowers. Other species of Cinquefoil (*Potentilla* spp.) are difficult to distinguish.

### **Monitoring and Rapid Response**

Hand-pulling or dig up, or by spot spraying of herbicides. The root crown (upper portion of the root system) must be removed or killed so that plants cannot resprout.

### **Credits**

The information provided in this factsheet was gathered from the Burke Museum of Natural History and Culture, Robert W. Freckmann Herbarium, USDA PLANTS Database and the U.S. Department of Agriculture, Forest Service. Individual species images that appear with a number in a black box are courtesy of the Bugwood.org network (<http://www.invasive.org>). Individual photo author credits may not be included due to the small display size of the images and subsequent difficulty of reading the provided text. All other images appear courtesy of Google (<http://images.google.com>).