

## Japanese stiltgrass

### *Microstegium vimineum*

#### Description

Introduced into the United States in Tennessee around 1919 and likely escaped as a result of its use as a packing material for porcelain.

#### Habit

Annual; grass, reaching 1 m (3.25 ft) in length. Lies on ground and roots at nodes along stems; upright branches look like individual plants.

#### Leaves

Alternate, pale green; asymmetrical; lanceolate; slightly pubescent; 5-8 cm (1.25-2 in) long, with a silvery line along the center of the blade.

#### Stems

Thin, weak roots form along the stem at nodes.

#### Flowers

Terminal racemes bear chasmogamous flowers,



Leslie J. Mehrhoff, University of Connecticut



David J. Moorhead  
University of Georgia



Nancy Loewenstein  
Auburn University

UGA2154034



Rebekah D. Wallace, University of Georgia

5426567



James H. Miller  
USDA Forest Service

058A0514368

while axillary racemes bear cleistogamous flowers.

### **Fruits and Seeds**

2.8-3.0 mm, ellipsoid achenes.

### **Habitat**

Native to Asia and Japan; shade tolerant/ does not tolerate full sun or standing water. Found in streams and riverbanks, moist forest, roadsides, ditches on moist soils.

### **Reproduction**

By seed. Very abundant which attach to animals, clothing; appears to be dispersed by deer. Seed persists in the seed bank for a long time.

### **Similar**

Native whitegrass (*Leersia virginica*).

### **Monitoring and Rapid Response**

Monitor shady paths, roadsides, moist sites in the fall; effectively controlled using any of the several readily available general use herbicides such as glyphosate.

### **Credits**

The Michigan Natural Features Inventory (MNFI) has partnered with MISIN to provide the information in this fact sheet. Species images and/or information were used with permission from "A Field Identification Guide to Invasive Plants in Michigan's Natural Communities" and "A Field Guide to Invasive Plants of Aquatic and Wetland Habitats for Michigan."