

## Spotted lanternfly

### *Lycorma delicatula*

#### Description

First detected in Pennsylvania in September of 2014.

#### Identification

Adults grow to about 1 inch long and 1/2 inch wide, and have large visually striking wings. Forewings are light brown with black spots at the front and a speckled band at the rear. Hind wings are scarlet with black spots at front and white and black bars at the rear. Yellow abdomen with black bars. Nymphs appear black with white spots and turn red before coming adults.

#### Habitat

Native to China, India, Japan, South Korea and Vietnam. Feeds on a variety of host plants including fruit trees, ornamental trees, woody trees and vines. Apples, birch, cheery, dogwood, grapes, Korean evodia, lilac, maple, poplar, stone fruits, and tree-of-heaven are among more than 70 species of host attacked.





## **Reproduction**

Eggs are laid on tree trunks in groups of 30-50, are yellowish brown in color, and covered with a gray, waxy coating prior to hatching. Hatch in early spring and early summer and nymphs begin feeding by sucking sap from young stems and leaves.

## **Impact**

Nymphs and adults cause damage when they feed, sucking sap from stems and leaves. Can reduce photosynthesis, weaken the plant and eventually contribute to the plant's death. Feeding can cause the plant to ooze or weep, resulting in a fermented odor. Insects excrete a large amount of fluid (honeydew) which promotes mold growth and attracts other insects.

## **Similar**

Hummingbird moths (*Hemaris* sp.), Tiger moths (*Arctiinae* sp.) and underwing moths (*Catocala* sp.) both share the bright and showy hind wing contrasting a duller fore wing. Poor flyer compared to native moths and insects, but is a strong, quick jumper.

## **Monitoring and Rapid Response**

Neonicotinoids, pyrethrins, and organophosphates are among effective chemical insecticides. Sticky traps at the base of trees and the parasitic wasp, *Anastatus orientalis* is reported to parasitize up to 69% of eggs in China.

## **Credits**

The information provided in this factsheet was gathered from University of California Agriculture and Natural Resources, Pennsylvania Dept. of Agriculture and USDA APHIS Pest Alert. Individual species images that appear with a number in a black box are courtesy of the Bugwood.org network (<https://www.invasive.org>). Individual photo author credits may not be included due to the small display

size of the images and subsequent difficulty of reading the provided text. All other images appear courtesy of Google (<https://images.google.com>).