

Cylindro

Cylindrospermopsis raciborskii

Description

Cylindro was likely introduced to the Great Lakes via ballast water. It can be transported to new water bodies by migrating waterfowl, recreational items, or natural water flow.

Habit

One of many species of blue-green algae composed of trichomes (chained filaments) that are solitary and straight. Trichomes range from 51?311 ?m in length and 1.7?4.2 ?m in width, heterocytes are 5?11 ?m by 2?5 ?m, and akinetes are 8?16 ?m by 2?5 ?m

Leaves

Stems

Flowers

Fruits and Seeds





Habitat

Native to South America. Found in low densities in many inland lakes and reservoirs surrounding the Great Lakes region.

Reproduction

Similar

Monitoring and Rapid Response

No known biological control. A mechanical system can be used to create artificial de-stratification to increase vertical mixing, introduce oxygen, and reduce internal nutrient loading. Can be inactivated by chlorine, ozone, and hydroxyl radical treatments. Copper-based algicides may inhibit the degradation.

Credits

The information provided in this factsheet was gathered from the Michigan Department of Natural Resources and the Great Lakes Aquatic Nonindigenous Information System. Individual species images that appear with a number in a black box are courtesy of the Bugwood.org network (<http://www.invasive.org>). Individual photo author credits may not be included due to the small display size of the images and subsequent difficulty of reading the provided text. All other images appear courtesy of Google (<http://images.google.com>).