

Pond water-starwort

Callitriche stagnalis

Description

Difficult to distinguish from other *Callitriche* species unless fruit is present; survey in late summer and fall. It can be distinguished by its fruits, which are almost round (orbicular).

Habit

Aquatic; perennial, delicate.

Leaves

Tiny, opposite, entire, forms vary, floating leaves spatulate, submergent leaves linear.

Stems

10-30 cm (4-12 in) in length, usually branched, rising to surface or sprawling.

Flowers

Tiny flowers lack sepals and petals and are located at the leaf bases on minute stalks, 2-4





tiny whitish bracts emerge from the flower base.

Fruits and Seeds

Small, located at the leaf bases. Four compartments, each containing one seed. Pond: Oval, 1.2-1.8 mm long, 1.2-1.7 mm wide, narrow margin all around (wing); bracts at base. Autumnal: 1.1 to 1.6 mm long, 1.2 to 1.8 mm wide, no margin, fruit without bracts.

Habitat

Native to Europe. Found in shallow waters of pond, marches or in protected areas in lakes or streams with slow-moving water.

Reproduction

By seed and clonally, dispersed by boats, vehicles, and animals-the latter both by fragments and ingested seed. Not all produce seeds.

Similar

Hard to distinguish from other water-starworts (*Callitriche* spp.) when not in fruit.

Monitoring and Rapid Response

Hand-pulling, cutting or raking making sure to remove all fragments. Clearing off boats and other water-related equipment will help prevent spreading.

Credits

The Michigan Natural Features Inventory (MNFI) has partnered with MISIN to provide the information in this fact sheet. Species images and/or information were used with permission from "A Field Identification Guide to Invasive Plants in Michigan's Natural Communities" and "A Field Guide to Invasive Plants of Aquatic and Wetland Habitats for Michigan."